

APPENDIX C. FLORISTIC BASELINE DATA

(Baseline Floristic Data for Tables 18, 19, and 20 in the Mitigation Success Criteria Section)

Appendix C. Plant Species by Habitat Type at the NVG Project Site

| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|--|----------------------------|---------------------|-------------------|------------------|
| <i>Vernal Pool (25 total, 17 native, 8 nonnative)</i> | | | | |
| <i>Anagallis arvensis</i> * | Scarlet Pimpernel | AH | FAC | Primulaceae |
| <i>Briza minor</i> * | Little Quakinggrass | AG | FACW- | Poaceae |
| <i>Callitrichia marginata</i> + | Winged Water-starwort | AH | OBL | Callitrichaceae |
| <i>Chamomilla suaveolens</i> * | Pineapple Weed | AH | FACU | Asteraceae |
| <i>Castilleja campestris</i> ssp. <i>campestris</i> | Field Owl's Clover | AH | OBL* | Orobanchaceae |
| <i>Deschampsia danthonioides</i> | Annual Hairgrass | AG | FACW | Poaceae |
| <i>Eleocharis macrostachya</i> | Creeping or Pale Spikerush | PG | OBL | Cyperaceae |
| <i>Epilobium ciliatum</i> | Northern Willow-herb | PH | FACW | Onagraceae |
| <i>Epilobium densiflorum</i> | Dense-flowered Willow-herb | AH | OBL | Onagraceae |
| <i>Eremocarpus setigerus</i> | Dove Weed | AH | - | Euphorbiaceae |
| <i>Gratiola ebracteata</i> | Bractless Hedgehyssop | AH | OBL | Scrophulariaceae |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> * | Mediterranean Barley | AG | FAC | Poaceae |
| <i>Juncus bufonius</i> | Common Toad Rush | AG | OBL | Juncaceae |
| <i>Lasthenia fremontii</i> | Fremont's Goldfields | A/PH | OBL | Asteraceae |
| <i>Lasthenia glaberrima</i> + | Smooth Goldfields | AH | OBL | Asteraceae |
| <i>Lolium multiflorum</i> * | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Lythrum hyssopifolium</i> * | Hyssop Loosestrife | AH | FACW | Lythraceae |
| <i>Navarretia leucocephala</i> + | Whitehead Navarretia | AH | OBL | Polemoniaceae |
| <i>Plagiobothrys stipitatus</i> + | Stalked Popcornflower | AH | OBL | Boraginaceae |
| <i>Poa annua</i> * | Annual Bluegrass | AG | FACW- | Poaceae |
| <i>Pseudognaphalium luteoalbum</i> * | Everlasting Cudweed | AH | FACW- | Asteraceae |
| <i>Psilocarphus brevissimus</i> + | Dwarf Woollyheads | AH | OBL | Asteraceae |
| <i>Ranunculus bonariensis</i> + | Carter's Buttercup | AH | OBL | Ranunculaceae |
| <i>Triteleia hyacinthina</i> | White Brodiaea | PH | FACW* | Liliaceae |
| <i>Veronica peregrina</i> | Neckweed | AH | OBL | Veronicaceae |
| <i>Vernal Pool Buffer (21 total, 4 native, 17 nonnative)</i> | | | | |
| <i>Anagallis arvensis</i> * | Scarlet Pimpernel | AH | FAC | Primulaceae |

³³ * = Introduced plant species that have become naturalized. + = Native species characteristic of the habitat. Scientific names of the plant species follow Hickman (1993) and Flora of North America Committee (2001-2007). Brackets [] indicate updated nomenclature.

³⁴ Habit definitions: AG = annual graminoid; AH = annual herb; AV = annual vine; F = Fern; PG = perennial graminoid; PH = perennial herb; PV = perennial vine; S = shrub; T = tree.

³⁵ WIS = Wetland Indicator Status. The following code definitions are according to Reed (1988):

OBL = obligate wetland species, occurs almost always in wetlands (>99% probability).

FACW = facultative wetland species, usually found in wetlands (67-99% probability).

FAC = facultative species, equally likely to occur in wetlands or nonwetlands (34-66% probability).

FACU = facultative upland species, usually found in nonwetlands (67-99% probability).

UPL = obligate upland species in this region (99% probability), occurs in wetlands in another region

NI = no indicator status has been assigned due to a lack of information.

+ or - symbols are modifiers that indicate greater or lesser affinity for wetland habitats.

* = tentative assignment to that indicator status by Reed (1988).

() Parentheses indicate a wetland status suggested by David L. Magney based on extensive field observations.

Daru – North Vineyard Greens Wetland Mitigation Plan

DMEC Project No.: 06-0112

August 2007



| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|---|----------------------------|---------------------|-------------------|----------------|
| <i>Avena barbata</i> * | Slender Wild Oat | A/PG | - | Poaceae |
| <i>Brassica rapa</i> * | Field Mustard | AH | - | Brassicaceae |
| <i>Briza minor</i> * | Little Quakinggrass | AG | FACW- | Poaceae |
| <i>Bromus diandrus</i> * | Ripgut Brome | AG | (FACU) | Poaceae |
| <i>Bromus hordeaceus</i> * | Soft Brome | AG | FACU- | Poaceae |
| <i>Bromus madritensis</i> ssp. <i>rubens</i> * | Red Brome | AG | NI | Poaceae |
| <i>Centaureum muhlenbergii</i> | Monterey Centaury | AH | FAC | Gentianaceae |
| <i>Chamomilla suaveolens</i> * | Pineapple Weed | AH | FACU | Asteraceae |
| <i>Deschampsia danthonioides</i> | Annual Hairgrass | AG | FACW | Poaceae |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> * | Mediterranean Barley | AG | FAC | Poaceae |
| <i>Hordeum murinum</i> * | Summer Barley | AG | NI | Poaceae |
| <i>Juncus capitatus</i> * | Leafybract Dwarf Rush | AG | FACU | Juncaceae |
| <i>Leontodon taraxacoides</i> * | Hawkbit | A/B/PH | FACU | Asteraceae |
| <i>Lepidium nitidum</i> | Common Peppergrass | AH | - | Brassicaceae |
| <i>Lolium multiflorum</i> * | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Lythrum hyssopifolium</i> * | Hyssop Loosestrife | AH | FACW | Lythraceae |
| <i>Plantago erecta</i> | California Plantain | AH | - | Plantaginaceae |
| <i>Poa annua</i> * | Annual Bluegrass | AG | FACW- | Poaceae |
| <i>Pseudognaphalium luteoalbum</i> * | Everlasting Cudweed | AH | FACW- | Asteraceae |
| <i>Vulpia bromoides</i> * | Brome Fescue | AG | FACW | Poaceae |
| <i>Seasonal Wetlands and Swales (26 total, 15 native, 11 nonnative)</i> | | | | |
| <i>Alisma lanceolatum</i> * | Lanceleaf Water Plantain | PH | OBL | Alismataceae |
| <i>Azolla filiculoides</i> | Pacific Mosquitofern | F | OBL | Azollaceae |
| <i>Crypsis schoenoides</i> * | Swamp Grass | AG | OBL | Poaceae |
| <i>Cyperus eragrostis</i> + | Umbrella-sedge | PG | FACW | Cyperaceae |
| <i>Deschampsia danthonioides</i> | Annual Hairgrass | AG | FACW | Poaceae |
| <i>Epilobium ciliatum</i> + | Northern Willow-herb | PH | FACW | Onagraceae |
| <i>Epilobium densiflorum</i> | Dense-flowered Willow-herb | AH | OBL | Onagraceae |
| <i>Epilobium pygmaeum</i> | Smooth Spike-primrose | AH | OBL | Onagraceae |
| <i>Glyceria declinata</i> * | Waxy Mannagrass | PG | - | Poaceae |
| <i>Gnaphalium palustre</i> + | Lowland Cudweed | AH | FACW | Asteraceae |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> * | Mediterranean Barley | AG | FAC | Poaceae |
| <i>Juncus balticus</i> + | Baltic Rush | PG | OBL | Juncaceae |
| <i>Juncus bufonius</i> | Common Toad Rush | AG | OBL | Juncaceae |
| <i>Juncus effusus</i> | Common Rush | PG | OBL | Juncaceae |
| <i>Juncus xiphioides</i> | Iris-leaved Rush | PG | OBL | Juncaceae |
| <i>Lolium multiflorum</i> * | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Mentha pulegium</i> * | Pennyroyal | PH | OBL | Lamiaceae |
| <i>Phyla nodiflora</i> | Turkey Tangle Fogfruit | PH | FACW | Verbenaceae |
| <i>Plagiobothrys stipitatus</i> + | Stalked Popcornflower | AH | OBL | Boraginaceae |
| <i>Polypogon monspeliensis</i> * | Rabbitsfoot Grass | AG | FACW+ | Poaceae |

Daru – North Vineyard Greens Wetland Mitigation Plan

DMEC Project No.: 06-0112

August 2007



| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|---|--------------------------|---------------------|-------------------|----------------|
| <i>Rumex acetosella</i> * ³⁶ | Common Sheep Sorrel | PH | FAC- | Polygonaceae |
| <i>Rumex crispus</i> * | Curly Dock | PH | FACW- | Polygonaceae |
| <i>Rumex pulcher</i> * | Fiddle Dock | PH | FAC+ | Polygonaceae |
| <i>Triteleia hyacinthina</i> | White Brodiaea | PH | FACW* | Liliaceae |
| <i>Veronica anagallis-aquatica</i> * | Water Speedwell | PH | OBL | Veronicaceae |
| <i>Veronica peregrina</i> | Neckweed | AH | OBL | Veronicaceae |
| <i>Seasonal Wetlands and Swales Buffer (23 total, 5 native, 18 nonnative)</i> | | | | |
| <i>Amaranthus retroflexus</i> + ³⁷ | Redroot Amaranth | AH | FACU | Amaranthaceae |
| <i>Anthemis cotula</i> * | Mayweed | AH | FACU | Asteraceae |
| <i>Asclepias fascicularis</i> + | Narrowleaf Milkweed | PH | FAC | Apocynaceae |
| <i>Bromus diandrus</i> * | Ripgut Brome | AG | (FACU) | Poaceae |
| <i>Bromus hordeaceus</i> * | Soft Brome | AG | FACU- | Poaceae |
| <i>Bromus madritensis</i> ssp. <i>rubens</i> * | Red Brome | AG | NI | Poaceae |
| <i>Chamomilla suaveolens</i> * | Pineapple Weed | AH | FACU | Asteraceae |
| <i>Chenopodium album</i> * | Lambsquarters | AH | FAC | Chenopodiaceae |
| <i>Festuca arundinacea</i> * | Tall Fescue | PG | FAC- | Poaceae |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> * | Mediterranean Barley | AG | FAC | Poaceae |
| <i>Lactuca serriola</i> * | Prickly Wild Lettuce | AH | FAC | Asteraceae |
| <i>Lepidium nitidum</i> + | Common Peppergrass | AH | - | Brassicaceae |
| <i>Lolium multiflorum</i> * | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Lotus corniculatus</i> * | Birdsfoot Trefoil | PH | FAC | Fabaceae |
| <i>Lythrum hyssopifolium</i> * | Hyssop Loosestrife | AH | FACW | Lythraceae |
| <i>Picris echioides</i> * | Bristly Ox-tongue | AH | (FAC) | Asteraceae |
| <i>Plantago erecta</i> + | California Plantain | AH | - | Plantaginaceae |
| <i>Plantago lanceolata</i> * | English Plantain | PH | FAC- | Plantaginaceae |
| <i>Polygonum arenastrum</i> * | Common Knotweed | AH | FAC | Polygonaceae |
| <i>Pseudognaphalium luteoalbum</i> * | Everlasting Cudweed | AH | FACW- | Asteraceae |
| <i>Ranunculus muricatus</i> * | Spinyfruit Buttercup | A/B/PH | FACW+ | Ranunculaceae |
| <i>Rumex crispus</i> * | Curly Dock | PH | FACW- | Polygonaceae |
| <i>Xanthium strumarium</i> + | Cocklebur | AH | FAC+ | Asteraceae |
| <i>Seasonal Marsh (17 total, 11 native, 6 nonnative)</i> | | | | |
| <i>Alisma lanceolatum</i> * | Lanceleaf Water Plantain | PH | OBL | Alismataceae |
| <i>Azolla filiculoides</i> | Pacific Mosquitofern | F | OBL | Azollaceae |
| <i>Crypsis schoenoides</i> * | Swamp Grass | AG | OBL | Poaceae |
| <i>Epilobium pygmaeum</i> | Smooth Spike-primrose | AH | OBL | Onagraceae |
| <i>Glyceria declinata</i> * | Waxy Mannagrass | PG | - | Poaceae |
| <i>Juncus effusus</i> + | Common Rush | PG | OBL | Juncaceae |
| <i>Juncus xiphioides</i> + | Iris-leaved Rush | PG | OBL | Juncaceae |
| <i>Lemna minuscula</i> | Least Duckweed | PH | OBL | Lemnaceae |
| <i>Ludwigia peploides</i> | Floating Water-primrose | PH | OBL | Onagraceae |
| <i>Mentha pulegium</i> * | Pennyroyal | PH | OBL | Lamiaceae |

Daru – North Vineyard Greens Wetland Mitigation Plan

DMEC Project No.: 06-0112

August 2007



| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|---|----------------------|---------------------|-------------------|----------------|
| <i>Polygonum hydropiperoides</i> | Swamp Smartweed | PH | OBL | Polygonaceae |
| <i>Polygonum punctatum</i> | Dotted Smartweed | A/PH | OBL | Polygonaceae |
| <i>Polypogon monspeliensis*</i> | Rabbitsfoot Grass | AG | FACW+ | Poaceae |
| <i>Ranunculus muricatus*</i> | Spinyfruit Buttercup | A/B/PH | FACW+ | Ranunculaceae |
| <i>Schoenoplectus [Scirpus] acutus+</i> | Hardstem Bulrush | PH | OBL | Cyperaceae |
| <i>Typha latifolia+</i> | Cattail | PH | OBL | Typhaceae |
| <i>Xanthium strumarium</i> | Cocklebur | AH | FAC+ | Asteraceae |
| <i>Seasonal Marsh Buffer (24 total, 5 native, 19 nonnative)</i> | | | | |
| <i>Anagallis arvensis*</i> | Scarlet Pimpernel | AH | FAC | Primulaceae |
| <i>Bromus diandrus*</i> | Ripgut Brome | AG | (FACU) | Poaceae |
| <i>Bromus hordeaceus*</i> | Soft Brome | AG | FACU- | Poaceae |
| <i>Bromus madritensis ssp. rubens*</i> | Red Brome | AG | NI | Poaceae |
| <i>Chamomilla suaveolens*</i> | Pineapple Weed | AH | FACU | Asteraceae |
| <i>Chenopodium album*</i> | Lambsquarters | AH | FAC | Chenopodiaceae |
| <i>Epilobium ciliatum+</i> | Northern Willow-herb | PH | FACW | Onagraceae |
| <i>Eryngium vaseyi+</i> | Coyotethistle | PH | FACW | Apiaceae |
| <i>Festuca arundinacea*</i> | Tall Fescue | PG | FAC- | Poaceae |
| <i>Lactuca serriola*</i> | Prickly Wild Lettuce | AH | FAC | Asteraceae |
| <i>Lepidium nitidum+</i> | Common Peppergrass | AH | - | Brassicaceae |
| <i>Lolium multiflorum*</i> | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Lotus corniculatus*</i> | Birdsfoot Trefoil | PH | FAC | Fabaceae |
| <i>Lythrum hyssopifolium*</i> | Hyssop Loosestrife | AH | FACW | Lythraceae |
| <i>Picris echioides*</i> | Bristly Ox-tongue | AH | (FAC) | Asteraceae |
| <i>Plantago erecta</i> | California Plantain | AH | - | Plantaginaceae |
| <i>Plantago lanceolata*</i> | English Plantain | PH | FAC- | Plantaginaceae |
| <i>Polygonum arenastrum*</i> | Common Knotweed | AH | FAC | Polygonaceae |
| <i>Polypogon monspeliensis*</i> | Rabbitsfoot Grass | AG | FACW+ | Poaceae |
| <i>Pseudognaphalium luteoalbum*</i> | Everlasting Cudweed | AH | FACW- | Asteraceae |
| <i>Rumex acetosella*</i> | Common Sheep Sorrel | PH | FAC- | Polygonaceae |
| <i>Rumex crispus*</i> | Curly Dock | PH | FACW- | Polygonaceae |
| <i>Rumex pulcher*</i> | Fiddle Dock | PH | FAC+ | Polygonaceae |
| <i>Xanthium strumarium+</i> | Cocklebur | AH | FAC+ | Asteraceae |
| <i>Upland Grassland (79 total, 18 native, 61 nonnative)</i> | | | | |
| <i>Aegilops triuncialis*</i> | Barbed Goatgrass | AG | - | Poaceae |
| <i>Aira caryophyllea*</i> | Silver Hairgrass | AG | - | Poaceae |
| <i>Amsinckia menziesii</i> | Rancher's Fire | AH | - | Boraginaceae |
| <i>Asclepias fascicularis</i> | Narrowleaf Milkweed | PH | FAC | Apocynaceae |
| <i>Avena barbata*</i> | Slender Wild Oat | A/PG | - | Poaceae |
| <i>Avena fatua*</i> | Wild Oat | AG | - | Poaceae |
| <i>Brachypodium distachyon*</i> | Purple False Brome | A/PG | - | Poaceae |
| <i>Brassica nigra*</i> | Black Mustard | AH | - | Brassicaceae |

Daru – North Vineyard Greens Wetland Mitigation Plan

DMEC Project No.: 06-0112

August 2007



| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|--|-------------------------|---------------------|-------------------|-----------------|
| <i>Brassica rapa</i> * | Field Mustard | AH | - | Brassicaceae |
| <i>Briza minor</i> * | Little Quakinggrass | AG | FACW- | Poaceae |
| <i>Brodiaea coronaria</i> | Harvest Brodiaea | PH | (FAC) | Liliaceae |
| <i>Bromus carinatus</i> | California Brome | AG | - | Poaceae |
| <i>Bromus diandrus</i> * | Ripgut Brome | AG | (FACU) | Poaceae |
| <i>Bromus hordeaceus</i> * | Soft Brome | AG | FACU- | Poaceae |
| <i>Bromus madritensis</i> ssp. <i>rubens</i> * | Red Brome | AG | NI | Poaceae |
| <i>Carduus pycnocephalus</i> * | Italian Thistle | AH | - | Asteraceae |
| <i>Castilleja attenuata</i> | Valley Tassels | AH | - | Orobanchaceae |
| <i>Centaurea solstitialis</i> * | Yellow Star-thistle | AH | - | Asteraceae |
| <i>Cerastium glomeratum</i> * | Mouse-ear Chickweed | AH | FACU | Caryophyllaceae |
| <i>Chamomilla suaveolens</i> * | Pineapple Weed | AH | FACU | Asteraceae |
| <i>Cichorium intybus</i> * | Chicory | PH | - | Asteraceae |
| <i>Cirsium vulgare</i> * | Bull Thistle | PH | FACU | Asteraceae |
| <i>Convolvulus arvensis</i> * | Bind Weed | PV | - | Convolvulaceae |
| <i>Crassula tillaea</i> * | Water Pygmy-weed | AH | NI* | Crassulaceae |
| <i>Cynodon dactylon</i> * | Bermuda Grass | PG | FAC | Poaceae |
| <i>Daucus carota</i> * | Queen Anne's Lace | PH | - | Apiaceae |
| <i>Epilobium brachycarpum</i> | Tall Annual Willow-herb | AH | UPL | Onagraceae |
| <i>Eremocarpus setigerus</i> | Dove Weed | AH | - | Euphorbiaceae |
| <i>Erodium botrys</i> * | Broadleaf Filaree | AH | - | Geraniaceae |
| <i>Erodium moschatum</i> * | Whitestem Filaree | AH | - | Geraniaceae |
| <i>Euphorbia spathulata</i> | Warty Spurge | AH | - | Euphorbiaceae |
| <i>Galium aparine</i> | Goose Grass | AH | FACU | Rubiaceae |
| <i>Geranium dissectum</i> * | Cut-leaved Geranium | AH | - | Geraniaceae |
| <i>Grindelia camporum</i> | Great Valley Gumplant | PH | FACU | Asteraceae |
| <i>Hemizonia fitchii</i> | Fitch's Tarweed | AH | - | Asteraceae |
| <i>Hirschfeldia incana</i> * | Summer Mustard | PH | - | Brassicaceae |
| <i>Holocarpha virgata</i> | Yellowflower Tarweed | AH | - | Asteraceae |
| <i>Hordeum marinum</i> ssp. <i>gussoneanum</i> * | Mediterranean Barley | AG | FAC | Poaceae |
| <i>Hordeum murinum</i> * | Summer Barley | AG | NI | Poaceae |
| <i>Hypochaeris glabra</i> * | Smooth Cat's-ear | AH | - | Asteraceae |
| <i>Kickxia elatine</i> * | Arrowleaf Fluvellin | AH | NI* | Veronicaceae |
| <i>Lactuca serriola</i> * | Prickly Wild Lettuce | AH | FAC | Asteraceae |
| <i>Lathyrus angulatus</i> * | Angled Pea | AV | - | Fabaceae |
| <i>Lepidium nitidum</i> | Common Peppergrass | AH | - | Brassicaceae |
| <i>Lolium multiflorum</i> * | Italian Ryegrass | AG | FAC* | Poaceae |
| <i>Lotus corniculatus</i> * | Birdsfoot Trefoil | PH | FAC | Fabaceae |
| <i>Lotus purshianus</i> | Spanish Clover | AH | UPL | Fabaceae |
| <i>Malva parviflora</i> * | Cheeseweed | AH | - | Malvaceae |
| <i>Medicago polymorpha</i> * | Burclover | AH | - | Fabaceae |

Daru – North Vineyard Greens Wetland Mitigation Plan

DMEC Project No.: 06-0112

August 2007



| Scientific Name ³³ | Common Name | Habit ³⁴ | WIS ³⁵ | Family |
|-------------------------------------|--------------------------|---------------------|-------------------|-----------------|
| <i>Paspalum dilatatum</i> * | Dallisgrass | PG | FAC | Poaceae |
| <i>Phalaris aquatica</i> * | Bulbous Canarygrass | PG | FAC+ | Poaceae |
| <i>Phytolacca americana</i> * | American Pokeweed | PH | NI | Phytolaccaceae |
| <i>Picris echioides</i> * | Bristly Ox-tongue | AH | (FAC) | Asteraceae |
| <i>Plantago erecta</i> | California Plantain | AH | - | Plantaginaceae |
| <i>Plantago lanceolata</i> * | English Plantain | PH | FAC- | Plantaginaceae |
| <i>Poa annua</i> * | Annual Bluegrass | AG | FACW- | Poaceae |
| <i>Polygonum arenastrum</i> * | Common Knotweed | AH | FAC | Polygonaceae |
| <i>Raphanus raphanistrum</i> * | Wild Radish | A/PH | - | Brassicaceae |
| <i>Raphanus sativus</i> * | Radish | A/BH | - | Brassicaceae |
| <i>Rumex acetosella</i> * | Common Sheep Sorrel | PH | FAC- | Polygonaceae |
| <i>Rumex pulcher</i> * | Fiddle Dock | PH | FAC+ | Polygonaceae |
| <i>Silene gallica</i> * | Windmill Pink | AH | - | Caryophyllaceae |
| <i>Silybum marianum</i> * | Milk Thistle | AH | - | Asteraceae |
| <i>Sonchus oleraceus</i> * | Common Sow-thistle | AH | NI* | Asteraceae |
| <i>Sorghum halepense</i> * | Johnsongrass | PG | FACU | Poaceae |
| <i>Spergularia rubra</i> * | Purple (Red) Sandspurrey | A/PH | FAC- | Caryophyllaceae |
| <i>Stellaria media</i> * | Common Chickweed | AH | FACU | Caryophyllaceae |
| <i>Taeniatherum caput-medusae</i> * | Medusahead | AG | - | Poaceae |
| <i>Tanacetum parthenium</i> * | Feverfew | PH | - | Asteraceae |
| <i>Taraxacum officinale</i> * | Dandelion | PH | FACU | Asteraceae |
| <i>Trichostema lanceolatum</i> | Vinegarweed | AH | - | Lamiaceae |
| <i>Trifolium dubium</i> * | Suckling Clover | AH | FACU* | Fabaceae |
| <i>Trifolium hirtum</i> * | Rose Clover | AH | - | Fabaceae |
| <i>Trifolium repens</i> * | White Clover | PH | FACU+ | Fabaceae |
| <i>Triteleia laxa</i> | Ithuriel's Spear | PG | - | Liliaceae |
| <i>Vicia sativa</i> * | Common Vetch | AH | FACU | Fabaceae |
| <i>Vicia villosa</i> * | Hairy Vetch | AH | - | Fabaceae |
| <i>Vulpia bromoides</i> * | Brome Fescue | AG | FACW | Poaceae |
| <i>Wyethia angustifolia</i> | California Compassplant | PH | FACU- | Asteraceae |